arXiv:1403.3201v1 [math.CA] 13 Mar 2014

The rate of convergence of some Riemann-Stieltjes sums

Adrian Holhoş

March 14, 2014

Abstract

We give the rate of convergence of some optimal lower Riemann-Stieltjes sums toward the integral.

1 Introduction

Let $[a,b]$ be a bounded closed interval. Let f,g be two functions defined on $[a,b]$. Consider an n-division Δ of $[a,b]$ defined by

$$\Delta\colon\ a=t_0<t_1<t_2<\cdots<t_n=b$$

and consider $\xi=(\xi_1,\xi_2,\ldots,\xi_n)$ such that $\xi_i\in[t_{i-1},t_i]$ for every $1\leq i\leq n$. The Riemann-Stieltjes sum is defined by

$$RS(f,g,\Delta,\xi)=\sum_{i=1}^{n}f(\xi_i)\cdot[g(t_i)-g(t_{i-1})].$$

The function f is said to be Riemann-Stieltjes integrable with respect to g if there is an $I\in\mathbb{R}$ with the property that for every $\varepsilon>0$ there exists $\delta>0$ such that for every division Δ of $[a,b]$ with mesh $\|\Delta\|=\max_{1\leq i\leq n}(t_i-t_{i-1})$ less than δ and every choice of the points ξ_i in $[t_{i-1},t_i]$ we have

$$|RS(f,g,\Delta,\xi)-I|<\varepsilon.$$

The number I is denoted $\int_a^b f(t)\,\mathrm{d}g(t)$ and is called the Riemann-Stieltjes integral of f with respect to g. When $g(x)=x$ we obtain the Riemann integrability.

Consider the lower Riemann-Stieltjes sum of a continuous function f on $[a,b]$

$$RS(f,g,\Delta,\min)=RS(f,g,\Delta,\xi),$$

where the points ξ_i are chosen such that $f(\xi_i)=\min_{t\in[a,b]}f(t)$. The set of all n-divisions of $[a,b]$ is compact and $\Delta\mapsto RS(f,g,\Delta,\min)$ is continuous, so there is an optimal n-division Δ_{opt} at which the lower Riemann-Stieltjes

sum is maximum. This optimal n-division may not be unique, but the sum $RS(f, g, \Delta_{opt}, \min)$ is unique.

In Theorem 6 we give the rate of approximation of the Riemann-Stieltjes integral by the optimal lower Riemann-Stieltjes sums, a result which generalizes Theorem 1.2 of [2].

2 Main results

We give next a generalization of a Lemma found in [1].

Lemma 1. Let $g\colon [a,b] \to \mathbb{R}$ be a non-negative continuous function which is not identically zero on any open subinterval of $[a,b]$ and let $h : [a,b] \to \mathbb{R}$ be a strictly positive and continuous function on $[a,b]$. For any positive integer n there exists a division of $[a,b]$:

$$t_0 = a < t_1 < t_2 < \cdots < t_{n-1} < t_n = b$$

such that the quantities

$$(t_i - t_{i-1}) \cdot \max_{t\in[t_{i-1},t_i]} g(t) \cdot \max_{t\in[t_{i-1},t_i]} h(t), \qquad 1 \le i \le n$$

are all equal to each other. Moreover, if J_n is the common value of all these quantities, then

$$\lim_{n\to\infty} nJ_n = \int_a^b g(t)h(t)\,\mathrm{d}t.$$

Proof. Parametrize the $(n-1)$ simplex σ by n-tuples $(u_1, u_2, \ldots, u_n)$, where $u_i \ge 0$ and $\sum_{i=1}^n u_i = 1$. Let this n-tuple correspond to the partition of $[a,b]$ given by

$$t_i = a + (b-a)\,(u_1 + u_2 + \cdots + u_i), \quad 1 \le i \le n, \text{ and } t_0 = a.$$

Let us define the function

$$\psi(u_1, u_2, \ldots, u_n) = (w_1, w_2, \ldots, w_n), \quad w_i = \frac{v_i}{\sum_{i=1}^n v_i},$$

where $v_i = (b-a)\,u_i \cdot \max_{t\in[t_{i-1},t_i]} g(t) \cdot \max_{t\in[t_{i-1},t_i]} h(t)$.

We have

$$\sum_{i=1}^n v_i = \sum_{i=1}^n (t_i - t_{i-1}) \max_{t\in[t_{i-1},t_i]} g(t) \cdot \max_{t\in[t_{i-1},t_i]} h(t)$$

is an upper Riemman sum for $\int_a^b g(t)h(t)\,\mathrm{d}t$ and $\sum_{i=1}^n v_i > 0$.

Since the maximum value of a continuous function over a closed interval depends continuously on the endpoints of that interval, ψ is a continuous function. Because $w_i = 0$ implies $u_i = 0$, ψ maps every face of σ into itself.

All this prove that ψ is surjective. So there exists $(u_1, u_2, \ldots, u_n)$ such that $\psi(u_1, u_2, \ldots, u_n) = (\frac{1}{n}, \frac{1}{n}, \ldots, \frac{1}{n})$. This proves the first part of our Lemma.

We have $nJ_n \leq (b-a)\,\|g\| \cdot \|h\|$. Let $\varepsilon > 0$ be given. From the continuity of g and h on $[a,b]$ there is a $\delta > 0$ so that $|t - t'| < \delta$ implies $|g(t) - g(t')| < \varepsilon$ and $|h(t) - h(t')| < \varepsilon$. We choose $n > \frac{(b-a)\|g\| \cdot \|h\|}{\varepsilon^2 \delta}$.

If $\max_{t \in [t_{i-1}, t_i]} g(t) \geq \varepsilon$ and $\max_{t \in [t_{i-1}, t_i]} h(t) \geq \varepsilon$ we have

$$J_n = (t_i - t_{i-1}) \cdot \max_{t \in [t_{i-1}, t_i]} g(t) \cdot \max_{t \in [t_{i-1}, t_i]} h(t) \geq \varepsilon^2 (t_i - t_{i-1}),$$

which proves that

$$t_i - t_{i-1} \leq \frac{J_n}{\varepsilon^2} = \frac{nJ_n}{n\varepsilon^2} \leq \frac{(b-a)\,\|g\| \cdot \|h\|}{n\varepsilon^2} < \delta.$$

This implies that the oscillations of g and of h over $[t_{i-1}, t_i]$ are at most ε. Considering η_i and ξ_i the points of maximum for g and h over the interval $[t_{i-1}, t_i]$ and applying the Mean Value Theorem for integrals we obtain

$$\begin{aligned}
\left| nJ_n - \int_a^b g(t)h(t)\,\mathrm{d}t \right| &= \left| \sum_{i=1}^{n} \left[g(\eta_i)h(\xi_i) - g(c_i)h(c_i) \right] (t_i - t_{i-1}) \right| \\
&\leq \sum_{i=1}^{n} \left(|g(\eta_i) - g(c_i)| \cdot |h(\xi_i)| + |g(c_i)| \cdot |h(\xi_i) - h(c_i)| \right) (t_i - t_{i-1}) \\
&\leq \varepsilon(\|h\| + \|g\|)(b-a).
\end{aligned}$$

This proves that nJ_n tends to $\int_a^b g(t)h(t)\,\mathrm{d}t$.

Consider now the case when $\max_{t \in [t_{i-1}, t_i]} g(t) < \varepsilon$ or $\max_{t \in [t_{i-1}, t_i]} h(t) < \varepsilon$. Suppose $g(t) < \varepsilon$ for every $t \in [t_{i-1}, t_i]$. The case when $\max_{t \in [t_{i-1}, t_i]} h(t) < \varepsilon$ can be analysed similarly. Because g is nonnegative we deduce also that the oscillation of g over the interval $[t_{i-1}, t_i]$ is at most ε. As we have done before nJ_n differs from the integral $\int_a^b g(t)h(t)\,\mathrm{d}t$ by less than $\varepsilon(b-a)3\,\|h\|$. □

Lemma 2. For every function $f \in C^1[a,b]$ and every $g \in C^1[a,b]$ with $g'(t) > 0$, for every $t \in [a,b]$, we have

$$\int_a^b f(t)\,\mathrm{d}g(t) - [g(b) - g(a)] \min_{t \in [a,b]} f(t) \leq \frac{1}{2}(b-a)^2\,\|f'\| \cdot \|g'\|\,.$$

Proof. Let $c \in [a,b]$ be the minimum point of f over $[a,b]$. We have

$$\begin{aligned}
\int_a^b f(t)\,\mathrm{d}g(t) - [g(b) - g(a)] \min_{t \in [a,b]} f(t) &= \int_a^b [f(t) - f(c)] g'(t)\,\mathrm{d}t \\
&\leq \|f'\| \cdot \|g'\| \cdot \int_a^b |t - c|\,\mathrm{d}t.
\end{aligned}$$

The proof is completed by using the inequality:

$$\int_a^b |t-c|\,\mathrm{d}t = \frac{(c-a)^2}{2} + \frac{(b-c)^2}{2} \le \frac{(b-a)^2}{2}.$$

□

Lemma 3. Consider f a function of class C^1 defined on $[a,b]$ with the derivative f' having a finite number of zeros. Let $g \in C^1[a,b]$ be a function with $g'(t) > 0$ for every t in $[a,b]$. Then

$$\limsup_{n\to\infty}\ n\left(\int_a^b f(t)\,\mathrm{d}g(t) - RS(f,g,\Delta_{opt},\min)\right) \le \frac{1}{2}\left(\int_a^b \sqrt{|f'(t)|\cdot g'(t)}\,\mathrm{d}t\right)^2.$$

Proof. We apply Lemma 1 to the functions $|f'(t)|^{\frac{1}{2}}$ and $|g'(t)|^{\frac{1}{2}}$ and obtain a division $\Delta' :\ a = t_0 < t_1 < t_2 < \cdots < t_n = b$ such that

$$J_n = (t_i - t_{i-1}) \cdot \max_{t\in[t_{i-1},t_i]} |f'(t)|^{\frac{1}{2}} \cdot \max_{t\in[t_{i-1},t_i]} |g'(t)|^{\frac{1}{2}},$$

has the same value for all values of $i \in \{1,2,\ldots,n\}$ and

$$\lim_{n\to\infty} nJ_n = \int_a^b \sqrt{|f'(t)|\cdot g'(t)}\,\mathrm{d}t.$$

Using Lemma 2 we obtain

$$\begin{aligned}\int_a^b f(t)\,\mathrm{d}g(t) - RS(f,g,\Delta',\min) &= \sum_{i=1}^n \left(\int_{t_{i-1}}^{t_i} f(t)\,\mathrm{d}g(t) - [g(t_i)-g(t_{i-1})] \min_{t\in[t_{i-1},t_i]} f(t)\right)\\ &\le \frac{1}{2}\sum_{i=1}^n (t_i - t_{i-1})^2 \cdot \max_{t\in[t_{i-1},t_i]} |f'(t)| \cdot \max_{t\in[t_{i-1},t_i]} |g'(t)|\end{aligned}$$

and finally

$$\begin{aligned}&\limsup_{n\to\infty}\ n\left(\int_a^b f(t)\,\mathrm{d}g(t) - RS(f,g,\Delta_{opt},\min)\right)\\ &\quad\le \limsup_{n\to\infty}\ n\left(\int_a^b f(t)\,\mathrm{d}g(t) - RS(f,g,\Delta',\min)\right)\\ &\quad\le \limsup_{n\to\infty} n\frac{1}{2}nJ_n^2 = \frac{1}{2}\lim_{n\to\infty}(nJ_n)^2\\ &\quad= \frac{1}{2}\left(\int_a^b \sqrt{|f'(t)|\cdot g'(t)}\,\mathrm{d}t\right)^2.\end{aligned}$$

□

Lemma 4. Consider f a function of class C^1 defined on $[a,b]$. Let $g \in C^1[a,b]$ be a function with $g'(t) > 0$ for every t in $[a,b]$. If $f'(t) \neq 0$ in a subinterval $[p,q]$ of $[a,b]$, then for every $\xi \in [p,q]$ we have

$$\left| \int_p^q f(t)\,\mathrm{d}g(t) - [g(q)-g(p)] \min_{t\in[p,q]} f(t) - \frac{1}{2}(q-p)^2 |f'(\xi)| g'(\xi) \right| \\ \leq \frac{1}{2}(q-p)^2 \cdot [\|g'\| \cdot \omega(f', q-p) + \|f'\| \cdot \omega(g', q-p)]$$

where $\omega(h,\delta)$ is the usual modulus of continuity of the function h.

Proof. Suppose $f' > 0$ on $[p,q]$. The case when the derivative of f is strictly negative on $[p,q]$ can be treated similarly. Because f is strictly increasing the minimum of f is attained in p. We have

$$\int_p^q f(t)\,\mathrm{d}g(t) - [g(q)-g(p)] \min_{t\in[p,q]} f(t) = \int_p^q [f(t)-f(p)] g'(t)\,\mathrm{d}t.$$

Applying the Mean Value Theorem for integrals twice we obtain

$$\begin{aligned} \int_p^q [f(t)-f(p)] g'(t)\,\mathrm{d}t &= g'(c) \cdot \int_p^q [f(t)-f(p)]\,\mathrm{d}t \\ &= g'(c) \cdot \int_p^q \int_p^t f'(u)\,\mathrm{d}u\,\mathrm{d}t \\ &= g'(c) \cdot \int_p^q f'(u)(q-u)\,\mathrm{d}u \\ &= g'(c) \cdot f'(d) \cdot \frac{(q-p)^2}{2}, \end{aligned}$$

for some $c, d \in (p,q)$. Because

$$|g'(c) \cdot f'(d) - f'(\xi) g'(\xi)| \leq \|g'\| \cdot \omega(f', q-p) + \|f'\| \cdot \omega(g', q-p)$$

the proof is complete. □

Lemma 5. Consider f a function of class C^1 defined on $[a,b]$ with the derivative f' having a finite number of zeros. Let $g \in C^1[a,b]$ be a function with $g'(t) > 0$ for every t in $[a,b]$. Then

$$\liminf_{n\to\infty} \; n \left(\int_a^b f(t)\,\mathrm{d}g(t) - RS(f,g,\Delta_{opt},\min) \right) \geq \frac{1}{2} \left(\int_a^b \sqrt{|f'(t)| \cdot g'(t)}\,\mathrm{d}t \right)^2.$$

Proof. We first prove that for any $\delta > 0$ there exists a positive integer r such that for any n-division Δ of $[a,b]$ the following inequality is true:

$$(n+r)^{\frac{1}{2}} \left(\int_a^b f(t)\,\mathrm{d}g(t) - RS(f,g,\Delta,\min) \right)^{\frac{1}{2}} \geq \frac{1}{\sqrt{2}} \int_a^b \sqrt{|f'(t)| g'(t)}\,\mathrm{d}t - \delta(b-a).$$

Since the function $x \mapsto x^{1/2}$ is uniformly continuous on $[0,\infty)$, there exists $\delta_1 > 0$ such that for any x and y in $[0,\infty)$ if $|x-y| < \delta_1$ then $|x^{1/2} - y^{1/2}| < \delta$. We take a subinterval $[p,q]$ of $[a,b]$ and suppose $f'(t) \neq 0$ in $[p,q]$. Because of the continuity of the derivatives of f and g there exists $\eta > 0$ such that if $q-p < \eta$ then $\frac{1}{2}\left[\|g'\| \cdot \omega(f', q-p) + \|f'\| \cdot \omega(g', q-p)\right] < \delta_1$. Using Lemma 4 we obtain

$$\left| \frac{\int_p^q f(t)\,\mathrm{d}g(t) - [g(q)-g(p)] \min_{t\in[p,q]} f(t)}{(q-p)^2} - \frac{1}{2}|f'(\xi)|g'(\xi)| \right| \leq \delta_1,$$

for any $\xi \in [p,q]$. Therefore, we have

$$\left| \frac{\left(\int_p^q f(t)\,\mathrm{d}g(t) - [g(q)-g(p)] \min_{t\in[p,q]} f(t)\right)^{\frac{1}{2}}}{q-p} - \frac{1}{\sqrt{2}}\sqrt{|f'(\xi)|g'(\xi)|} \right| \leq \delta,$$

which is equivalent with

$$\left| \left(\int_p^q f(t)\,\mathrm{d}g(t) - [g(q)-g(p)] \min_{t\in[p,q]} f(t)\right)^{\frac{1}{2}} - \frac{1}{\sqrt{2}}\sqrt{|f'(\xi)|g'(\xi)|}(q-p) \right| \leq \delta(q-p).$$

Since f' is uniformly continuous on $[a,b]$, for the above $\delta > 0$ there exists $\zeta > 0$ such that $|x-y| < \zeta$ implies $|f'(x) - f'(y)| < \delta^2 / \|g'\|$. We denote by Z the zero set of f':

$$Z = \{\, t \in [a,b] \,|\, f'(t) = 0 \,\}$$

and define the ζ-neighborhood Z_ζ of Z by

$$Z_\zeta = \{\, u \in [a,b] \,|\, \exists\, t \in Z :\ |t-u| < \zeta \,\}.$$

Then for any $t \in Z_\zeta$ we have $g'(t)|f'(t)| < \delta^2$ and f' is not equal to 0 on the complement of Z_ζ. By the definition of Z_ζ and the properties of f' we can see that Z_ζ is a disjoint union of finitely many intervals (by choosing ζ small enough). We denote by r_1 the number of all endpoints of the intervals of Z_ζ. For $\eta > 0$ obtained above we take a positive integer r_2 satisfying $r_2 \geq (b-a)/\eta$ and set $r = r_1 + r_2$. For any n-division Δ of $[a,b]$ we can add at most r_2 points to Δ such that the mesh of the new division is less than or equal to η. Moreover we add the endpoints of all the intervals of Z_ζ and denote the new division by

$$\Delta' : t_0 = a < t_1 < \ldots, < t_m = b.$$

By the definition of Δ' we have $m \leq n + r$ and $t_i - t_{i-1} \leq \eta$. Each interval $[t_{i-1}, t_i]$ satisfies $[t_{i-1}, t_i] \subset \overline{Z_\zeta}$ or $[t_{i-1}, t_i] \subset [a,b] \setminus Z_\zeta$. In both cases we can take $c_i \in [t_{i-1}, t_i]$ satisfying

$$\int_{t_{i-1}}^{t_i} \sqrt{|f'(t)| \cdot g'(t)}\,\mathrm{d}t = \sqrt{|f'(c_i)| \cdot g'(c_i)}(t_i - t_{i-1}).$$

In the case $[t_{i-1}, t_i] \subset \overline{Z_\zeta}$ we have

$$\begin{aligned}
&\frac{1}{\sqrt{2}}\sqrt{|f'(c_i)| \cdot g'(c_i)}(t_i - t_{i-1}) \le \frac{1}{\sqrt{2}}\delta(t_i - t_{i-1}) \\
&\quad \le \left(\int_{t_i}^{t_{i-1}} f(t)\,\mathrm{d}g(t) - [g(t_i) - g(t_{i-1})] \min_{t\in[t_{i-1},t_i]} f(t)\right)^{\frac{1}{2}} + \delta(t_i - t_{i-1}).
\end{aligned}$$

In the case $[t_{i-1}, t_i] \subset [a, b] \setminus Z_\zeta$, f' is not equal to 0 in $[t_{i-1}, t_i]$, so

$$\begin{aligned}
&\frac{1}{\sqrt{2}}\sqrt{|f'(c_i)| \cdot g'(c_i)}(t_i - t_{i-1}) \\
&\quad \le \left(\int_{t_i}^{t_{i-1}} f(t)\,\mathrm{d}g(t) - [g(t_i) - g(t_{i-1})] \min_{t\in[t_{i-1},t_i]} f(t)\right)^{\frac{1}{2}} + \delta(t_i - t_{i-1}).
\end{aligned}$$

Adding all these inequalities for $i = 1, 2, \ldots, m$ we get

$$\begin{aligned}
&\frac{1}{\sqrt{2}}\int_a^b \sqrt{|f'(t)| \cdot g'(t)}\,\mathrm{d}t \\
&\quad \le \sum_{i=1}^{m}\left(\int_{t_i}^{t_{i-1}} f(t)\,\mathrm{d}g(t) - [g(t_i) - g(t_{i-1})] \min_{t\in[t_{i-1},t_i]} f(t)\right)^{\frac{1}{2}} + \delta(b - a).
\end{aligned}$$

Applying the Cauchy-Schwarz inequality to the first term of the right-hand side of the above inequality we obtain

$$\begin{aligned}
&\sum_{i=1}^{m}\left(\int_{t_i}^{t_{i-1}} f(t)\,\mathrm{d}g(t) - [g(t_i) - g(t_{i-1})] \min_{t\in[t_{i-1},t_i]} f(t)\right)^{\frac{1}{2}} \\
&\quad \le m^{\frac{1}{2}}\left(\sum_{i=1}^{m}\left(\int_{t_i}^{t_{i-1}} f(t)\,\mathrm{d}g(t) - [g(t_i) - g(t_{i-1})] \min_{t\in[t_{i-1},t_i]} f(t)\right)\right)^{\frac{1}{2}} \\
&\quad = m^{\frac{1}{2}}\left(\int_a^b f(t)\,\mathrm{d}g(t) - RS(f, g, \Delta', \min)\right)^{\frac{1}{2}}.
\end{aligned}$$

From these we have

$$\frac{1}{\sqrt{2}}\int_a^b \sqrt{|f'(t)| \cdot g'(t)}\,\mathrm{d}t \le m^{\frac{1}{2}}\left(\int_a^b f(t)\,\mathrm{d}g(t) - RS(f, g, \Delta', \min)\right)^{\frac{1}{2}} + \delta(b - a).$$

Because

$$[g(c) - g(b)] \min_{t\in[b,c]} f(t) + [g(b) - g(a)] \min_{t\in[a,b]} f(t) \ge [g(c) - g(a)] \min_{t\in[a,c]} f(t),$$

for every $a < b < c$, we obtain

$$\int_a^b f(t)\,\mathrm{d}g(t) - RS(f, g, \Delta', \min) \le \int_a^b f(t)\,\mathrm{d}g(t) - RS(f, g, \Delta, \min).$$

This estimate and $m \leq n + r$ imply

$$\frac{1}{\sqrt{2}} \int_a^b \sqrt{|f'(t)|g'(t)}\,\mathrm{d}t \leq (n+r)^{\frac{1}{2}} \left(\int_a^b f(t)\,\mathrm{d}g(t) - RS(f,g,\Delta,\min) \right)^{\frac{1}{2}} + \delta(b-a).$$

Now, let us prove Lemma 5. From the continuity of the function $x \mapsto x^2$ in $x_0 = \frac{1}{\sqrt{2}} \int_a^b \sqrt{|f'(t)|g'(t)}\,\mathrm{d}t$, for any $\varepsilon > 0$ there exists $\xi > 0$ such that if $x_0 - x \leq \xi$ we have $x_0^2 - x^2 \leq \frac{\varepsilon}{2}$. We take $\delta > 0$ which satisfies $\delta(b-a) < \xi$. We can apply the result obtained above and get a positive integer r such that for any n-division Δ of $[a,b]$

$$\begin{aligned} \xi &\geq \delta(b-a) \\ &\geq \frac{1}{\sqrt{2}} \int_a^b \sqrt{|f'(t)|g'(t)}\,\mathrm{d}t - (n+r)^{\frac{1}{2}} \left(\int_a^b f(t)\,\mathrm{d}g(t) - RS(f,g,\Delta,\min) \right)^{\frac{1}{2}}, \end{aligned}$$

which implies

$$\frac{\varepsilon}{2} \geq \frac{1}{2} \left(\int_a^b \sqrt{|f'(t)|g'(t)}\,\mathrm{d}t \right)^2 - (n+r) \left(\int_a^b f(t)\,\mathrm{d}g(t) - RS(f,g,\Delta,\min) \right).$$

We can substitute the optimal division Δ_{opt} for Δ in the above inequality and get

$$(n+r) \left(\int_a^b f(t)\,\mathrm{d}g(t) - RS(f,g,\Delta_{opt},\min) \right) \geq \frac{1}{2} \left(\int_a^b \sqrt{|f'(t)|g'(t)}\,\mathrm{d}t \right)^2 - \frac{\varepsilon}{2}.$$

Since

$$\lim_{n\to\infty} \left(\int_a^b f(t)\,\mathrm{d}g(t) - RS(f,g,\Delta_{opt},\min) \right) = 0,$$

we can choose a positive integer N such that for $n \geq N$ the inequality

$$0 \leq r \left(\int_a^b f(t)\,\mathrm{d}g(t) - RS(f,g,\Delta_{opt},\min) \right) \leq \frac{\varepsilon}{2}$$

holds. Thus

$$n \left(\int_a^b f(t)\,\mathrm{d}g(t) - RS(f,g,\Delta_{opt},\min) \right) \geq \frac{1}{2} \left(\int_a^b \sqrt{|f'(t)|g'(t)}\,\mathrm{d}t \right)^2 - \varepsilon,$$

for every $n \geq N$. This completes the proof. □

Theorem 6. *Consider f a function of class C^1 defined on $[a,b]$ with the derivative f' having a finite number of zeros. Let $g \in C^1[a,b]$ be a function with $g'(t) > 0$ for every t in $[a,b]$. Then*

$$\lim_{n\to\infty} n \left(\int_a^b f(t)\,\mathrm{d}g(t) - RS(f,g,\Delta_{opt},\min) \right) = \frac{1}{2} \left(\int_a^b \sqrt{|f'(t)| \cdot g'(t)}\,\mathrm{d}t \right)^2.$$

Proof. The result follows from the inequalities of Lemma 3 and 5. □

References

[1] A. M. Gleason, *A curvature formula*, Amer. J. Math., **101** (1979), 86–93.

[2] H. Tasaki, *Convergence rates of approximate sums of Riemann integrals*, J. Approx. Theory, **161**, 2 (2009), 477–490.